arXiv:1209.2692v1 [math.NA] 12 Sep 2012

Exact regularity of symmetric univariate subdivision schemes

Michael S. Floater* and Georg Muntingh†

September 13, 2012

Abstract

In this paper we review and refine a technique of Rioul to determine the Hölder regularity of a large class of symmetric subdivision schemes from the spectral radius of a single matrix. These schemes include those of Dubuc and Deslauriers, their dual versions, and more generally the pseudo-spline and dual pseudo-spline schemes. The technique also applies to convex combinations of these, and therefore to several 'tension parameter' schemes.

MSC: 65D10, 26A16
Keywords: Subdivision, Hölder regularity, pseudo-splines.

1 Introduction

Subdivision is a recursive method for generating curves, surfaces and other geometric objects. Rather than having a complete description of the object of interest at hand, subdivision generates the object by repeatedly refining its description starting from a coarse set of control points. Since subdivision schemes are often easy to implement and very flexible, they provide a powerful tool for modelling geometry. However, analyzing their smoothness,

---

*Centre of Mathematics for Applications, Department of Informatics, University of Oslo, PO Box 1053, Blindern, 0316 Oslo, Norway, *email: michaelf@ifi.uio.no*

†Centre of Mathematics for Applications, Department of Mathematics, University of Oslo, PO Box 1053, Blindern, 0316 Oslo, Norway, *email: georgmu@math.uio.no*

or regularity, can be difficult. The purpose of this paper is to review and refine a method proposed by Rioul [18] to determine the Hölder regularity of a surprisingly large class of subdivision schemes from the spectral radius of a single matrix. A joint spectral radius analysis is not required.

Consider the scheme

$$f_{j+1,k} = \sum_{\ell} a_{k-2\ell} f_{j,\ell}, \tag{1}$$

with finitely supported mask $\mathbf{a} = (a_k)_{k\in\mathbb{Z}}$, and coefficients $a_k \in \mathbb{R}$, acting on the initial data $f_{0,k} \in \mathbb{R}$, $k \in \mathbb{Z}$. At each subdivision level $j \geq 0$, let f_j be the piecewise linear function with value $f_{j,k}$ at the point $2^{-j}k$. The scheme is then *convergent* if it has a pointwise limit $f := \lim_{j\to\infty} f_j$. We will assume that only a finite number of the initial data $f_{0,k}$ are non-zero, in which case f has compact support. In the special case of the cardinal data $f_{0,k} = \delta_{k,0}$, the support of the limit f will be the interval $[K, L]$ if $a_K, a_L \neq 0$ and $a_k = 0$ for all $k < K$ and $k > L$. We note that shifting the a_k merely shifts f: if $\tilde{f}$ is the limit of the scheme with shifted coefficients $\tilde{a}_k := a_{k+1}$ then $\tilde{f}(x) = f(x+1)$.

The Laurent polynomial

$$A(z) = \sum_{k} a_k z^k \tag{2}$$

is the *symbol* of the scheme. We note that multiplying or dividing $A(z)$ by any power of z results in the symbol of the scheme with shifted coefficients, and therefore has no effect on the regularity of f: if for example $\tilde{a}_k := a_{k+1}$ then $\tilde{A}(z) = z^{-1}A(z)$. It has been shown, for example by Dyn and Levin [13], that a necessary condition for convergence of (1) is that

$$\sum_{k} a_{2k} = \sum_{k} a_{2k+1} = 1, \tag{3}$$

and so we will make this assumption. This condition can be expressed in terms of the symbol as

$$A(-1) = 0 \qquad \text{and} \qquad A(1) = 2. \tag{4}$$

Let us now suppose, after shifting the coefficients a_k as necessary, that $A(z)$ admits the factorization

$$A(z) = 2^{-r}(1+z)^{r+1}B(z) \tag{5}$$

for some $r \geq 0$, where $\mathbf{b}$, the masking corresponding to B, is *symmetric* about b_0, i.e., $b_k = b_{-k}$. Then the Fourier transform of $\mathbf{b}$,

$$\widehat{\mathbf{b}}(\xi) := B(e^{-i\xi}), \qquad \xi \in \mathbb{R}, \tag{6}$$

which is periodic with period 2π, is also real.

Rioul showed in [18] that a *lower bound* on the Hölder regularity of (1) can be determined from the spectral radius of a *single matrix* if $\widehat{\mathbf{b}} \geq 0$, i.e., if $\widehat{\mathbf{b}}(\xi) \geq 0$ for all $\xi \in [-\pi, \pi]$. A surprisingly large class of schemes are of this type. For example, one can easily check that they include all the pseudo-spline and dual pseudo-spline schemes, from the explicit formulas for $A(z)$ in [4, 8, 9, 7].

Rioul further showed that in the special case that the scheme (1) is interpolatory, the lower bound is optimal. We will show more: that the lower bound is optimal whenever the cardinal function of the scheme has ℓ^∞-stable integer translates, which includes the interpolatory schemes as a special case. Using a characterization of such stability due to Jia and Micchelli [17], this leads us to conclude that the lower bound is optimal under the slightly stricter condition that $\widehat{\mathbf{b}} > 0$. Such schemes include again all the pseudo-spline and dual pseudo-spline schemes.

To be more specific, recall that a function such as f has *Hölder regularity* α, $0 < \alpha < 1$, written $f \in C^\alpha$, if

$$|f(x) - f(y)| \leq C|x - y|^\alpha$$

for all $x, y \in \mathbb{R}$, and we write $f \in C^{q+\alpha}$ for $q \in \mathbb{N}_0$, $0 < \alpha < 1$, if $f \in C^q$, i.e., f is q times continuously differentiable, and $f^{(q)} \in C^\alpha$. Correspondingly, we shall say that the *scheme* (1) has Hölder regularity γ for some real $\gamma \geq 0$, if, for $\beta < \gamma$, $f \in C^\beta$ for all initial data, and, for $\beta > \gamma$, $f \notin C^\beta$ for some initial data.

By the assumption of the symmetry of $\mathbf{b}$, it must have the form

$$\mathbf{b} = (\ldots, 0, b_p, \ldots, b_1, b_0, b_1, \ldots, b_p, 0, \ldots), \qquad b_p \neq 0, \tag{7}$$

for some $p \geq 0$, in which case

$$\widehat{\mathbf{b}}(\xi) = b_0 + 2\sum_{k=1}^{p} b_k \cos(k\xi). \tag{8}$$

Let $M_{\mathbf{b}}$ be the $p \times p$ matrix defined by

$$(M_{\mathbf{b}})_{i+1,j+1} = \begin{cases} b_i & \text{if } j = 0; \\ b_{|i-2j|} + b_{i+2j} & \text{otherwise,} \end{cases} \tag{9}$$

so that the first few examples of $M_{\mathbf{b}}$, with $p = 1, 2, 3, 4$, are

$$\begin{bmatrix} b_0 \end{bmatrix}, \quad \begin{bmatrix} b_0 & 2b_2 \\ b_1 & b_1 \end{bmatrix}, \quad \begin{bmatrix} b_0 & 2b_2 & 0 \\ b_1 & b_1 + b_3 & b_3 \\ b_2 & b_0 & b_2 \end{bmatrix}, \quad \begin{bmatrix} b_0 & 2b_2 & 2b_4 & 0 \\ b_1 & b_1 + b_3 & b_3 & 0 \\ b_2 & b_0 + b_4 & b_2 & b_4 \\ b_3 & b_1 & b_1 & b_3 \end{bmatrix}. \tag{10}$$

Let ρ be the spectral radius of $M_{\mathbf{b}}$. Our main theorem is

Theorem 1. *If* $\widehat{\mathbf{b}} \geq 0$ *and* $\rho > 1/2$*, a lower bound for the Hölder regularity of* (1) *is* $r - \log_2(\rho)$*. If* $\widehat{\mathbf{b}} > 0$ *this bound is optimal.*

In practice one can take the largest possible r in (5), so that the matrix $M_{\mathbf{b}}$ is relatively small and has relatively large spectral radius. As an example, consider the quintic Dubuc-Deslauriers scheme with mask

$$\mathbf{a} = \frac{1}{256}(\ldots, 0, 3, 0, -25, 0, 150, 256, 150, 0, -25, 0, 3, 0, \ldots). \tag{11}$$

There is a factorization (5) up to $r = 5$, in which case one finds

$$\mathbf{b} = (b_{-2}, b_{-1}, b_0, b_1, b_2) = \frac{1}{8}(3, -18, 38, -18, 3) \tag{12}$$

and

$$M_{\mathbf{b}} = \frac{1}{8}\begin{bmatrix} 38 & 6 \\ -18 & -18 \end{bmatrix}. \tag{13}$$

Writing $s = \sin^2(\xi/2)$ yields

$$\widehat{\mathbf{b}}(\xi) = \frac{1}{8}(38 - 36\cos\xi + 6\cos 2\xi) = 1 + 3s + 6s^2 > 0 \tag{14}$$

for any $\xi \in \mathbb{R}$. Since $M_{\mathbf{b}}$ has the two eigenvalues $9/2$ and -2, Theorem 1 implies that this scheme has regularity

$$5 - \log_2(9/2) \approx 2.8301.$$

The remainder of the paper is organized as follows. In the next three sections, we recall some notation and results on the convergence and regularity of (1), and their relation to the growth rate of the 'derived scheme' defined by $B(z)$. The following three sections explain how to estimate this growth rate in the case that $\widehat{\mathbf{b}} \geq 0$ and the section after that shows that this estimate is sharp if $\widehat{\mathbf{b}} > 0$, proving Theorem 1. Finally, using the formulas of [4, 8, 9, 7], we apply the theorem to compute the Hölder regularity of the pseudo-spline and dual pseudo-spline subdivision schemes, and some convex combinations of these.

2 Regularity

The regularity of f is related to the behaviour of the divided differences of the scheme, and their differences, defined recursively as $f_{j,k}^{[0]} = f_{j,k}$ and $g_{j,k}^{[0]} = f_{j,k} - f_{j,k-1}$, and for $s \geq 1$,

$$f_{j,k}^{[s]} = \frac{2^j}{s} g_{j,k}^{[s-1]}, \qquad g_{j,k}^{[s]} = f_{j,k}^{[s]} - f_{j,k-1}^{[s]}. \tag{15}$$

If $A(z)$ is divisible by $(1+z)^{r+1}$ for some $r \geq 0$, as in (5), then there is a scheme for the $f_{j,k}^{[s]}$ and $g_{j,k}^{[s]}$ for $s = 0, 1, \ldots, r$. For such s, if we define their associated Laurent polynomials as

$$F_j^{[s]}(z) = \sum_k f_{j,k}^{[s]} z^k, \qquad G_j^{[s]}(z) = \sum_k g_{j,k}^{[s]} z^k,$$

then

$$F_{j+1}^{[s]}(z) = A^{[s]}(z) F_j^{[s]}(z^2), \tag{16}$$

and

$$G_{j+1}^{[s]}(z) = B^{[s]}(z) G_j^{[s]}(z^2), \tag{17}$$

where

$$A^{[s]}(z) = \frac{2^s}{(1+z)^s} A(z),$$

and

$$B^{[s]}(z) = \frac{2^s}{(1+z)^{s+1}} A(z),$$

from which we obtain the derived schemes

$$f_{j+1,k}^{[s]} = \sum_{\ell} a_{k-2\ell}^{[s]} f_{j,\ell}^{[s]}, \tag{18}$$

and

$$g_{j+1,k}^{[s]} = \sum_{\ell} b_{k-2\ell}^{[s]} g_{j,\ell}^{[s]}. \tag{19}$$

Under this divisibility condition, it can be shown [13] that a sufficient condition for f to be in C^r is that

$$|g_{j,k}^{[r]}| \leq C\lambda^j, \tag{20}$$

for some constants C and $\lambda < 1$, independent of j and k. Moreover, if $1/2 < \lambda < 1$, then $f \in C^{r-\log_2(\lambda)}$.

3 Reduction procedure

How can we use (20) in the case that it holds with $\lambda \geq 1$? Then we do not know whether $f \in C^r$, but if $r \geq 1$ we can use the 'reduction procedure' of Daubechies, Guskov, and Sweldens [2] to obtain information about lower order derivatives. Although the procedure was shown to work for interpolatory schemes in [2], it also applies to the more general scheme (1).

Lemma 1. *Suppose that $A(z)$ is divisible by $(1+z)^{r+1}$ for some $r \geq 1$. If (20) holds with $\lambda > 1$ then*

$$|g_{j,k}^{[r-1]}| \leq D_1 2^{-j} \lambda^j, \tag{21}$$

while if it holds with $\lambda = 1$,

$$|g_{j,k}^{[r-1]}| \leq (D_2 + D_3 j) 2^{-j}, \tag{22}$$

for constants D_1, D_2, D_3.

Proof. By the divisibility assumption, $A^{[r]}(-1) = 0$, and by the assumption that $A(1) = 2$ in (4), it also follows that $A^{[r]}(1) = 2$. Therefore,

$$\sum_k a_{2k}^{[r]} = \sum_k a_{2k+1}^{[r]} = 1, \tag{23}$$

and from (18) using summation by parts there is a constant C_1 such that

$$|f_{j+1,2k+s}^{[r]} - f_{j,k}^{[r]}| \leq C_1 \max_k |g_{j,k}^{[r]}|, \qquad s = 0, 1.$$

So, for any $j \geq 1$, if we represent any $k \in \mathbb{Z}$ in binary form as $k = k_j$, where

$$k_\ell = 2k_{\ell-1} + s_\ell, \qquad \ell = j, j-1, \ldots, 1,$$

for some $k_0 \in \mathbb{Z}$ and $s_1, \ldots, s_j \in \{0, 1\}$, then

$$|f_{j,k}^{[r]} - f_{0,k_0}^{[r]}| \leq \sum_{\ell=1}^{j} |f_{\ell,k_\ell}^{[r]} - f_{\ell-1,k_{\ell-1}}^{[r]}| \leq C_1 C(1 + \lambda + \cdots + \lambda^{j-1}).$$

Hence,

$$|f_{j,k}^{[r]}| \leq C_2 + C_1 C(1 + \lambda + \cdots + \lambda^{j-1}),$$

and since

$$g_{j,k}^{[r-1]} = 2^{-j} r f_{j,k}^{[r]},$$

this gives the result in the two cases $\lambda > 1$ and $\lambda = 1$. □

By applying this procedure recursively, it follows that if (20) holds for any λ with $1/2 < \lambda < 2^r$ then $f \in C^{r - \log_2 \lambda}$ if $\log_2 \lambda$ is not an integer, and $f \in C^{r - \log_2 \lambda - \epsilon}$ for any small $\epsilon > 0$ if $\log_2 \lambda$ is an integer.

4 Rioul's method

With r in (5) now fixed, we have $B(z) = B^{[r]}(z)$. If we also write $b_k = b_k^{[r]}$, $g_{j,k} = g_{j,k}^{[r]}$, and $G_j(z) = G_j^{[r]}$, the derived scheme for the $g_{j,k}^{[r]}$ can be expressed as

$$g_{j+1,k} = \sum_\ell b_{k-2\ell} g_{j,\ell} \tag{24}$$

or in terms of Laurent polynomials as

$$G_{j+1}(z) = B(z) G_j(z^2). \tag{25}$$

Rioul showed that if $B(z)$ has the form (7) and if $\widehat{\mathbf{b}} \geq 0$, (20) holds with λ the spectral radius of a single matrix whose elements are formed from the coefficients b_k. Note that if $p = 0$ we must have $b_0 = 1$ and so we can take

$\lambda = 1$. In fact in this case the scheme (1) is the B-spline scheme of degree r and this merely confirms the well-known fact that the limit f, being a spline of degree r, belongs to C^β for any $\beta < r$. Thus, we assume from now on that $p \geq 1$. Iterating (25) gives

$$G_j(z) = B_j(z) G_0(z^{2^j}), \tag{26}$$

where

$$B_j(z) := B(z)B(z^2) \cdots B(z^{2^{j-1}}). \tag{27}$$

But then

$$B_{j+1}(z) = B(z)B_j(z^2), \tag{28}$$

and so $B_j(z)$ is the Laurent polynomial of the data $b_{j,k}$, where $b_{0,k} = \delta_{k,0}$ and

$$b_{j+1,k} = \sum_\ell b_{k-2\ell} b_{j,\ell}. \tag{29}$$

In particular, $b_{1,k} = b_k$. Since (26) can be written as

$$g_{j,k} = \sum_\ell b_{j,k-2^j\ell} g_{0,\ell}, \tag{30}$$

it follows that

$$|g_{j,k}| \leq \max_r |b_{j,r}| \sum_\ell |g_{0,\ell}|,$$

and so (20) holds if there is some constant C' such that

$$\max_k |b_{j,k}| \leq C' \lambda^j. \tag{31}$$

By induction on j, the values $b_{j,k}$ are zero whenever $k < -p_j$ or $k > p_j$, where $p_j := (2^j - 1)p$.

Lemma 2 (Rioul)**.** *If $B(z)$ in (5) has the form (7) and $\widehat{\mathbf{b}} \geq 0$ then*

$$\max_k |b_{j,k}| = |b_{j,0}| \qquad \textit{for all } j \geq 0.$$

Proof. Since

$$\widehat{\mathbf{b}}_j(\xi) := \sum_{k=-p_j}^{p_j} b_{j,k} e^{-ik\xi}$$

is a Fourier series, we have

$$b_{j,k} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \widehat{\mathbf{b}}_j(\xi) e^{ik\xi} \, d\xi.$$

Therefore,

$$|b_{j,k}| \le \frac{1}{2\pi} \int_{-\pi}^{\pi} |\widehat{\mathbf{b}}_j(\xi)| \, d\xi.$$

By the symmetry of the b_k, it follows from (29) by induction on j that the $b_{j,k}$ are symmetric for all j, i.e., $b_{j,-k} = b_{j,k}$. Therefore, $\widehat{\mathbf{b}}_j$ is real, and by induction on j from (28), $\widehat{\mathbf{b}}_j(\xi) > 0$ for all ξ. It follows that

$$|b_{j,k}| \le \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} \widehat{\mathbf{b}}_j(\xi) \, d\xi \right| = |b_{j,0}|. \qquad \Box$$

5 A folded matrix

Under the assumption of Lemma 2 it follows from (31) that (20) holds if

$$|b_{j,0}| \le C' \lambda^j \tag{32}$$

for some constant C'. Thus we only need to estimate the grow rate of the central coefficient $b_{j,0}$ as $j \to \infty$. Since $b_{j,-k} = b_{j,k}$, one way to determine this rate is to study the behaviour of the vectors of coefficients

$$\mathbf{b}_j = (b_{j,0}, b_{j,1}, \ldots, b_{j,p-1})^T,$$

since they are self-generating and include $b_{j,0}$. Indeed, from (29),

$$b_{j+1,k} = b_k b_{j,0} + \sum_{\ell \ge 1} (b_{k-2\ell} + b_{k+2\ell}) b_{j,\ell},$$

and using the fact that $b_{-k} = b_k$ implies

$$b_{j+1,k} = b_k b_{j,0} + \sum_{\ell \ge 1} (b_{|k-2\ell|} + b_{k+2\ell}) b_{j,\ell},$$

and it follows that $\mathbf{b}_{j+1} = M \mathbf{b}_j$, where M is the $p \times p$ matrix $M_{\mathbf{b}}$ defined in (9). Therefore, with $\mathbf{e} = (1, 0, \ldots, 0)^T \in \mathbb{R}^p$,

$$|b_{j,0}| = \|M^j \mathbf{e}\|_\infty \le \|M^j\|_\infty \sim \rho(M)^j, \qquad \text{as } j \to \infty.$$

where $\rho(M)$ is the spectral radius of M. Thus (20) holds with $\lambda = \rho(M)$.

6 An alternative folded matrix

Rioul obtained the growth rate of $b_{j,0}$ in a different way, working from an alternative to equation (28). From (27) there is an alternative recursion:

$$B_{j+1}(z) = B(z^{2^j})B_j(z),$$

and therefore

$$b_{j+1,k} = \sum_{\ell} b_\ell b_{j,k-2^j\ell}. \tag{33}$$

This means that the subset of coefficients $b_{j,k}$ whose indices increase in steps of 2^j, rather than 1, i.e, $c_{j,k} := b_{j,2^jk}$, satisfy the equation

$$c_{j+1,k} = \sum_{\ell} b_\ell c_{j,2k-\ell} = \sum_{\ell} b_{2k-\ell} c_{j,\ell}. \tag{34}$$

Since $b_{j,0} = c_{j,0}$, we can obtain the growth rate of $b_{j,0}$ by studying the behaviour of the vectors

$$\mathbf{c}_j = (c_{j,0}, c_{j,1}, \ldots, c_{j,p-1})^T,$$

instead of the $\mathbf{b}_j$. Similar to (29), equation (34) simplifies since the mask $\mathbf{b}$ is symmetric, for then $c_{j,-\ell} = c_{j,\ell}$, and so

$$c_{j+1,k} = b_{2k}c_{j,0} + \sum_{\ell \geq 1}(b_{2k-\ell} + b_{2k+\ell})c_{j,\ell},$$

and again because $b_{-k} = b_k$,

$$c_{j+1,k} = b_{2k}c_{j,0} + \sum_{\ell \geq 1}(b_{|2k-\ell|} + b_{2k+\ell})c_{j,\ell}.$$

Thus $\mathbf{c}_{j+1} = N\mathbf{c}_j$, where N is the $p \times p$ matrix defined by

$$(N)_{i+1,j+1} = \begin{cases} b_{2i} & \text{if } j = 0; \\ b_{|2i-j|} + b_{2i+j} & \text{otherwise,} \end{cases} \tag{35}$$

The first few examples of N, with $p = 1, 2, 3, 4$, are

$$\begin{bmatrix} b_0 \end{bmatrix}, \quad \begin{bmatrix} b_0 & 2b_1 \\ b_2 & b_1 \end{bmatrix}, \quad \begin{bmatrix} b_0 & 2b_1 & 2b_2 \\ b_2 & b_1 + b_3 & b_0 \\ 0 & b_3 & b_2 \end{bmatrix}, \quad \begin{bmatrix} b_0 & 2b_1 & 2b_2 & 2b_3 \\ b_2 & b_1 + b_3 & b_0 + b_4 & b_1 \\ b_4 & b_3 & b_2 & b_1 \\ 0 & 0 & b_4 & b_3 \end{bmatrix}.$$

Therefore,

$$|b_{j,0}| = |c_{j,0}| = \|N^j \mathbf{e}\|_\infty = \|N^j\|_\infty \sim \rho(N)^j, \qquad \text{as } j \to \infty.$$

Thus we can instead take $\lambda = \rho(N)$. However, because the transpose N^T is similar to M, one has $\rho(N) = \rho(M)$.

7 Optimality

Let ϕ denote the cardinal function of the scheme (1), i.e., its limit when the initial data is the cardinal data $f_{0,k} = \delta_{k,0}$. Then the limit of the general scheme can be expressed as the linear combination

$$f(x) = \sum_\ell f_{0,\ell}\phi(x - \ell).$$

Following Jia and Micchelli [17], we shall say that ϕ *has ℓ^∞-stable integer translates* if there is some constant $K > 0$ such that for any sequence $\mathbf{c} = (c_\ell)_\ell$ in $\ell^\infty(\mathbb{Z})$,

$$\|\sum_\ell c_\ell \phi(\cdot - \ell)\|_{L^\infty(\mathbb{R})} \geq K\|\mathbf{c}\|_{\ell^\infty(\mathbb{Z})}. \tag{36}$$

Lemma 3. *Suppose ϕ has ℓ^∞-stable integer translates and f has regularity $q + \alpha$ for some $q \in \mathbb{N}_0$ and $0 < \alpha < 1$. Then for any integer $r \geq q$, there is a constant C such that*

$$|g^{[r]}_{j,k}| \leq C 2^{j(r-q-\alpha)}. \tag{37}$$

Proof. As is well known, see e.g. the review by Dyn and Levin [13], ϕ satisfies the two-scale difference equation

$$\phi(x) = \sum_k a_k \phi(2x - k), \tag{38}$$

and therefore, for any $j \geq 0$,

$$f(x) = \sum_\ell f_{j,\ell}\phi(2^j x - \ell). \tag{39}$$

We can use this equation to relate any divided difference of f of the form

$$\tilde{f}^{[q]}_{j,y} := [2^{-j}(y - q), 2^{-j}(y - q + 1), \ldots, 2^{-j}(y)]f,$$

for $y \in \mathbb{R}$, to the divided differences of the scheme. Putting $x = 2^{-j}(y-k)$ in (39) gives

$$f\big(2^{-j}(y-k)\big) = \sum_{\ell} f_{j,\ell-k}\phi(y-\ell),$$

and, using the cases $k = 0, 1, \ldots, q$, and the linearity of divided differences,

$$\tilde{f}^{[q]}_{j,y} = \sum_{\ell} f^{[q]}_{j,\ell}\phi(y-\ell).$$

Similarly, if

$$\tilde{g}^{[q]}_{j,y} := \tilde{f}^{[q]}_{j,y} - \tilde{f}^{[q]}_{j,y-1},$$

then

$$\tilde{g}^{[q]}_{j,y} = \sum_{\ell} g^{[q]}_{j,\ell}\phi(y-\ell).$$

Recalling that f has compact support, if f has regularity $q+\alpha$, there is some $C > 0$ such that for any $\xi_0, \xi_1 \in \mathbb{R}$,

$$|f^{(q)}(\xi_1) - f^{(q)}(\xi_0)| \le C|\xi_1 - \xi_0|^{\alpha},$$

and, by a standard property of divided differences, for each j and y,

$$|\tilde{g}^{[q]}_{j,y}| = |f^{(q)}(\xi_1) - f^{(q)}(\xi_0)|/q!,$$

for $\xi_0, \xi_1 \in \big(2^{-j}(y-q-1), 2^{-j}(y)\big)$. Therefore, for any y,

$$|\tilde{g}^{[q]}_{j,y}| \le C' 2^{-j\alpha},$$

where

$$C' = C(q+1)^{\alpha}/q!.$$

Therefore,

$$\|\sum_{\ell} g^{[q]}_{j,\ell}\phi(\cdot - \ell)\|_{L^\infty(\mathbb{R})} \le C' 2^{-j\alpha},$$

and by (36) it follows that for any $\ell \in \mathbb{Z}$,

$$|g^{[q]}_{j,\ell}| \le K^{-1} C' 2^{-j\alpha}.$$

Finally, by applying the divided difference definitions (15) recursively, we obtain (37). □

It follows from this lemma that as long as ϕ has ℓ^∞-stable integer translates, the lower bound on the Hölder regularity of the scheme that we obtained from the spectral radius $\rho = \rho(M)$ in Section 5 is optimal. To see this, suppose that ρ corresponds to a real eigenvalue of M. After taking the initial data

$$(g_{0,0}^{[r]}, \dots, g_{0,p}^{[r]})^T$$

in the derived scheme (24) as the corresponding eigenvector, extending by $g_{0,-k}^{[r]} = g_{0,k}^{[r]}$ yields a sequence of divided differences $g_{j,k}^{[r]}$ that grows at precisely the rate ρ^j, i.e.,

$$|g_{j,k}^{[r]}| \sim \rho^j. \tag{40}$$

In the other case that ρ corresponds to a pair of complex conjugate eigenvalues, we again obtain the growth rate (40) by taking the real part of one of the corresponding eigenvectors. Suppose now that $f \in C^{r-\log_2 \rho+\epsilon}$ for some small $\epsilon > 0$. Writing the exponent as

$$r - \log_2 \rho + \epsilon = q + \alpha,$$

for $q \in \mathbb{N}_0$ and $0 < \alpha < 1$, if $\rho > 1/2$, we have $q \leq r$, and the lemma then implies that

$$|g_{j,k}^{[r]}| \leq C 2^{j(\log_2 \rho - \epsilon)} = C\rho^j 2^{-j\epsilon},$$

which contradicts (40).

It remains to consider when ϕ has ℓ^∞-stable integer translates. If the scheme (1) is interpolatory, in the sense that $a_{2k} = \delta_{k,0}$, then $\phi(k) = \delta_{k,0}$ and so the stability condition (36) holds with $K = 1$. Thus, for the Dubuc-Deslauriers schemes the spectral radius of M yields the optimal regularity. Rioul showed this and tabulated the regularities of these schemes in [18].

To show stability in the general case, we apply some results by Jia and Micchelli [17]. We denote the (continuous) Fourier transform of ϕ by

$$\widehat{\phi}(\xi) = \int_{\mathbb{R}} \phi(x) e^{-i\xi x}\, dx, \qquad \xi \in \mathbb{R}.$$

Since the scheme (1) has constant precision,

$$\sum_{\ell} \phi(x - \ell) = 1, \qquad x \in \mathbb{R},$$

and, as shown by Jia and Micchelli [17, Theorem 2.4], $\widehat{\phi}(0) = 1$. Since the Fourier transform of (38) is

$$\widehat{\phi}(\xi) = 2^{-1}\widehat{\mathbf{a}}(\xi/2)\widehat{\phi}(\xi/2),$$

it follows that

$$\widehat{\phi}(\xi) = \prod_{j=1}^{\infty} \big(2^{-1}\widehat{\mathbf{a}}(\xi/2^j)\big).$$

A sufficient condition [17, Theorem 3.5] for ϕ to have ℓ^∞-stable integer translates is that

$$\sup_{\ell\in\mathbb{Z}} |\widehat{\phi}(\xi + 2\pi\ell)| > 0, \qquad \text{for all } \xi \in \mathbb{R}. \tag{41}$$

Consider then again the case that the derived scheme (24) holds. Then

$$\widehat{\mathbf{a}}(\xi) = 2\cos^{r+1}(\xi/2)\widehat{\mathbf{b}}(\xi),$$

where, since $\widehat{\mathbf{a}}(0) = 2$ under the assumption of convergence, $\widehat{\mathbf{b}}(0) = 1$. In the B-spline scheme of degree r we have $\widehat{\mathbf{b}}(z) = 1$, in which case we can write the symbol as $A_r(z) = (1+z)^{r+1}/2^r$. The cardinal function ϕ_r is the B-spline of degree r centred at 0, and we have

$$\widehat{\phi}_r(\xi) = \prod_{j=1}^{\infty} \cos^{r+1}\left(\xi/2^{j+1}\right) = \left(\frac{\sin(\xi/2)}{\xi/2}\right)^{r+1}.$$

It then follows that

$$\widehat{\phi}(\xi) = \widehat{\phi}_r(\xi) \prod_{j=1}^{\infty} \widehat{\mathbf{b}}\big(\xi/2^j\big).$$

Since the condition (41) holds for the B-spline ϕ_r we deduce that ϕ has ℓ^∞-stable integer translates if $\widehat{\mathbf{b}}(\xi) > 0$ for all $\xi \in \mathbb{R}$. This completes the proof of Theorem 1.

This idea of comparing with B-splines, which are known to be stable, was used by Dong and Shen [8, Lemma 2.2] to show that all pseudo-splines are stable.

8 Pseudo-splines and dual pseudo-splines

It turns out that a surprisingly large class of subdivision schemes that appear in the literature satisfy the conditions of Theorem 1, including the pseudo-spline and dual pseudo-spline schemes.

Consider the family of *primal schemes* with symbols

$$A_{m,l}(z) = 2\sigma^m(z) \sum_{k=0}^{l} \binom{m-1+k}{k} \delta^k(z), \qquad m \geq 1, \quad l \geq 0, \tag{42}$$

where

$$\sigma(z) = \frac{(1+z)^2}{4z}, \qquad \delta(z) = -\frac{(1-z)^2}{4z}.$$

It is shown by Dong and Shen [8] that these are the symbols of the pseudo-splines of type II, which were introduced by Daubechies, Han, Ron, and Shen [4] and studied systematically by Dong and Shen [9]. The $2m$-point Dubuc-Deslauriers schemes form the subfamily with symbols $A_{m,m-1}$. Other choices of l and m lead to other known schemes. Dividing $A_{m,l}$ by $2\sigma^m$ gives us the symbol for the shifted derived scheme,

$$B_{m,l}(z) = B_{m,l}^{[2m-1]}(z) = \sum_{k=0}^{l} \binom{m-1+k}{k} \delta^k(z) = \sum_{i=-l}^{l} b_{m,l,i} z^i, \tag{43}$$

with

$$b_{m,l,i} = (-1)^i \sum_{k=|i|}^{l} \binom{m-1+k}{k} \binom{2k}{i+k} 2^{-2k}. \tag{44}$$

A family of schemes analogous to the Dubuc-Deslauriers schemes was introduced by Dyn, Floater, and Hormann [11] and generalized by Dyn, Hormann, Sabin, and Shen [7] to a family of *dual schemes* with symbols

$$\tilde{A}_{m,l}(z) = \frac{1+z}{z} \sigma^m(z) \sum_{k=0}^{l} \binom{m-1/2+k}{k} \delta^k(z), \qquad m \geq 1, \quad l \geq 0. \tag{45}$$

Dividing $\tilde{A}_{m,l}(z)$ by $\frac{1+z}{z}\sigma^m(z)$ gives us the symbol for the shifted derived scheme,

$$\tilde{B}_{m,l}(z) = \tilde{B}_{m,l}^{[2m]}(z) = \sum_{k=0}^{l} \binom{m-1/2+k}{k} \delta^k(z) = \sum_{i=-l}^{l} \tilde{b}_{m,l,i} z^i, \tag{46}$$

with

$$\tilde{b}_{m,l,i} = (-1)^i \sum_{k=|i|}^{l} \binom{m - 1/2 + k}{k} \binom{2k}{i+k} 2^{-2k}.$$

For any $m \geq 1$ and $l \geq 0$, since

$$\delta\big(e^{i\xi}\big) = \sin^2(\xi/2) \geq 0, \qquad \xi \in [-\pi, \pi],$$

we see that $\widehat{\mathbf{b}}_{m,l} > 0$ and $\widehat{\tilde{\mathbf{b}}}_{m,l} > 0$ and so Theorem 1 can be applied.

Example 1. *Consider the primal schemes with* $l = 1$*. From (43–44) we find that* M *is a* 1×1 *matrix with spectral radius* $b_{m,1,0} = 1 + m/2 > 1/2$*. We conclude that the primal scheme with symbol* $A_{m,1}$ *has Hölder regularity* $2m - \log_2(m+2)$*, which was also shown by Hormann and Sabin [16].*

Example 2. *For the primal scheme with* $(m, l) = (4, 3)$*, one finds* $B_{4,3}(z) = b_{-3}z^{-3} + \cdots + b_3 z^3$*, with*

$$(b_{-3}, \dots, b_3) = \frac{1}{16}(-5, 40, -131, 208, -131, 40, -5).$$

The regularity of the original scheme with symbol $A_{4,3}$ *is then* $7 - \log_2(\rho) \approx 3.55113$*, with* $\rho \approx 10.91976$ *the largest zero (in absolute value) of the polynomial*

$$\det(M - \lambda I) = \det \begin{bmatrix} b_0 - \lambda & 2b_2 & 0 \\ b_1 & b_1 + b_3 - \lambda & b_3 \\ b_2 & b_0 & b_2 - \lambda \end{bmatrix} = -\lambda^3 + 7\lambda^2 + \frac{217}{4}\lambda - 125.$$

Similarly, one can, for any integers $m \geq 1$ and $l \geq 0$, compute the Hölder regularities of the schemes with symbols $A_{m,l}$ and $\tilde{A}_{m,l}$ as the $\log_2$ of algebraic numbers of degree at most l. For any integers $1 \leq l < m \leq 8$, Tables 1 and 2 contain Hölder regularities for the primal and dual schemes, showing that the first four decimals of the lower bounds presented by Dong, Dyn, and Hormann [6] are, in fact, significant.

Moreover, since

$$\sigma\big(e^{i\xi}\big) = \cos^2(\xi/2) \geq 0, \qquad \xi \in [-\pi, \pi],$$

any convex combination of the schemes $A_{m,l}$, i.e., the schemes with symbols

$$(1 - w)A_{m_1,l_1}(z) + wA_{m_2,l_2}(z), \qquad w \in [0, 1],$$

	$l=1$	$l=2$	$l=3$	$l=4$	$l=5$	$l=6$	$l=7$
$m=2$	2						
$m=3$	3.67807	2.83007					
$m=4$	5.41504	4.34379	3.55113				
$m=5$	7.19265	5.92502	4.96207	4.19357			
$m=6$	9	7.55781	6.43997	5.53250	4.77675		
$m=7$	10.83007	9.23111	7.97187	6.93577	6.06273	5.31732	
$m=8$	12.67807	10.93702	9.54804	8.39272	7.41006	6.56398	5.82944

Table 1: Hölder regularities rounded to five decimals for the primal schemes $A_{m,l}$ with $1 \leq l < m \leq 8$.

	$l=1$	$l=2$	$l=3$	$l=4$	$l=5$	$l=6$	$l=7$
$m=2$	2.83007						
$m=3$	4.54057	3.57723					
$m=4$	6.29956	5.12711	4.24726				
$m=5$	8.09311	6.73575	5.69355	4.85423			
$m=6$	9.91254	8.38994	7.19984	6.22682	5.41143		
$m=7$	11.75207	10.08039	8.75493	7.65811	6.72934	5.93283	
$m=8$	13.60768	11.80033	10.35034	9.13861	8.10385	7.20968	6.43070

Table 2: Hölder regularities rounded to five decimals for the dual schemes $\tilde{A}_{m,l}$ with $1 \leq l < m \leq 8$.

have a derived scheme satisfying the conditions of Theorem 1. The same is true for the dual schemes $\tilde{A}_{m,l}$.

Figure 1, for instance, shows Hölder regularities of convex combinations of the primal and dual Dubuc-Deslauriers schemes. Similarly, as remarked by Dyn et al. [7], for $L \geq 1$, the symbols of the schemes $S_{2L}(w)$ of Choi, Lee, Lee, and Yoon [1] are affine combinations of $A_{L,L-1}$ and $A_{L+1,L-1}$, while the symbols of their schemes $S_{2L-1}(w)$ are affine combinations of the duals $\tilde{A}_{L,L-1}$ and $\tilde{A}_{L-1,L-1}$. When the combinations are convex, Theorem 1 can be applied.

References

[1] S. W. Choi, B.-G. Lee, Y. J. Lee, J. Yoon, Stationary subdivision schemes reproducing polynomials, Comput. Aided Geom. Design **23**



Figure 1: Hölder regularities of the primal (left) and dual (right) Dubuc-Deslauriers $2m$-point schemes versus m, interpolated by the Hölder regularities of the corresponding tension parameter schemes.

(2006), no. 4, 351–360.

[2] I. Daubechies, I. Guskov, and W. Sweldens, Regularity of irregular subdivision, Constr. Approx. **15** (1999), 381–426.

[3] I. Daubechies, I. Guskov, and W. Sweldens, Commutation for irregular subdivision, Constr. Approx. **17** (2001), 479–513.

[4] I. Daubechies, B. Han, A. Ron, Z. Shen, Framelets: MRA-based constructions of wavelet frames, Appl. Comput. Harmon. Anal. **14** (2003), no. 1, 1–46.

[5] G. Deslauriers and S. Dubuc, Symmetric iterative interpolation processes, Constr. Approx. **5** (1989), 49–68.

[6] B. Dong, N. Dyn, and K. Hormann, Properties of dual pseudo-splines, Appl. Comp. Harm. Anal. **29** (2010), 104–110.

[7] N. Dyn, K. Hormann, M. A. Sabin, and Z. Shen, Polynomial reproduction by symmetric subdivision schemes, J. Approx. Theory **155** (2008), 28–42.

[8] B. Dong and Z. Shen, Linear independence of pseudo-splines, Proc. Amer. Math. Soc. **134** (2006), 2685–2694.

[9] B. Dong and Z. Shen, Pseudo-splines, wavelets and framelets, Appl. Comput. Harmon. Anal. **22** (2007), no. 1, 78–104.

[10] S. Dubuc, Interpolation through an iterative scheme, J. Math. Anal. Appl. **114** (1986), 185–204.

[11] N. Dyn, M. S. Floater, and K. Hormann, A C^2 four-point subdivision scheme with fourth order accuracy and its extensions. Mathematical methods for curves and surfaces: Tromsø 2004, 145–156.

[12] N. Dyn, M. S. Floater, and K. Hormann, Four-point curve subdivision based on iterated chordal and centripetal parameterizations, Comput. Aided Geom. Design. **26** (2009), 279–286.

[13] N. Dyn and D. Levin, Subdivision schemes in geometric modelling, Acta Numerica **11** (2002), 73–144.

[14] M. S. Floater, A piecewise polynomial approach to analyzing interpolatory subdivision, J. Approx. Theory **163** (2011), 1547–1563.

[15] J. Hechler, B. Mossner, and U. Reif, C^1-continuity of the generalized four-point scheme, Linear Algebra and its Applications, **430** (2009), 3019–3029.

[16] K. Hormann, M. A. Sabin, A family of subdivision schemes with cubic precision, Comput. Aided Geom. Design **25** (2008), no. 1, 41–52.

[17] R. Q. Jia, C. A. Micchelli, Using the refinement equations for the construction of pre-wavelets. II. Powers of two, Curves and surfaces: Chamonix-Mont-Blanc 1990, 209–246.

[18] O. Rioul, Simple regularity criteria for subdivision schemes, SIAM J. Math. Anal. **23** (1992), 1544–1576.

[19] M. J. Shensa, The discrete wavelet transform: wedding the à trous and mallat algorithms, IEEE Transactions on Signal Processing, Vol. 40. No. 10, (October 1992), 2464–2482.